

The nest and eggs of Red-winged Wood-rail *Aramides calopterus* in the foothills of north-east Ecuador

Jarol F.Vaca B., Harold F. Greeney, Rudolphe A. Gelis, Caroline Dingle, Neils Krabbe and Mike Tidwell

Received 10 September 2005; final revision received 20 February 2006

Cotinga 26(2006): 13–14

En la reserva de la comunidad de Mushullacta, que se encuentra dentro de la zona de amortiguamiento del Parque Nacional Sumaco-Galeras, a 1.175 msnm encontramos un nido de *Aramides calopterus* con tres huevos de color crema con manchas de color café y violeta claro. El nido se encontró entre dos pequeñas laderas con un pantano permanente y descansaba entre dos árboles con raíces zancudas; una palmera (*Iriatea deltoidea*) y otro árbol no identificado. Estimamos que el periodo de incubación sería de 24 días.



Figure 1. Adult Red-winged Wood-rail *Aramides calopterus* incubating three eggs, Mushullacta community reserve, Napo, Ecuador, 5 April 2005 (Murray Cooper)

The genus *Aramides* numbers seven medium to large rails (the wood-rails⁴). The genus is largely confined to South America, with only two species reaching Central America. Several are extremely poorly known and one (Brown Wood-rail *A. wolffi*) is considered Vulnerable¹. One of the least known, Red-winged Wood-rail *A. calopterus*, has a peculiar distribution, with apparently disjunct populations in the eastern foothills and lowlands of Ecuador, north-central and south-east Peru, and Amazonian



Figure 2. Nest and complete clutch of Red-winged Wood-rail *Aramides calopterus* Mushullacta community reserve, Napo, Ecuador, 4 April 2005 (Harold F. Greeney)

Brazil⁴. Nothing has been published concerning its natural history, even the voice being unknown.

Here we present data on the nest and eggs of *A. calopterus* from the foothills of north-east Ecuador. In April 2005, we made observations at 1,175 m elevation at a campsite (00°22'S 78°08'W) within a community-owned reserve, Mushullacta, Napo province, Ecuador, adjacent to Napo-Galeras National Park. For additional images of the nest, eggs and adults see Greeney².

Nest

The nest was discovered on 3 April 2005, at 18h15, when an adult was flushed and a nest with three eggs discovered. The nest was a shallow cup of dead leaves and sticks, supported at 1.1 m by the converging root masses of two small (25-cm dbh) trees; *Iriartea deltoidea* Palmae and an unknown dicot (Fig. 1). The nest was loosely constructed, with material seemingly piled into the niche provided by the roots. It measured roughly 22–25 cm in diameter and 13–14 cm tall outside. The cup was unlined and measured c.15 cm in diameter by 6 cm deep.

Eggs

The eggs were off-white to beige with dense splotches and spots of dark brown, red-brown and lavender (Fig. 2). At 07h30 on 4 April they measured 46.6 × 35.4 mm, 47.9 × 35.7 mm and 46.9 × 35.8 mm, and weighed 31.93, 32.87 and 32.52 g, respectively. Seven days later, at 18h00, they weighed 31.21, 31.93 and 31.71 g. This represents a mean mass-loss rate of 0.11 g/24 h or 0.34% per day. Using this mean, and an estimated incubation period of 24 days, we calculate a mean 8.2% loss of mass during incubation.

Additional observations and discussion

At 05h30 on 4 April there was no adult on the nest and the eggs were cold and wet, making us suspect that no adult spent the night at the nest. A strong light held behind the eggs revealed no visible embryonic development. We believe, therefore, that the nest was found shortly after the clutch was completed. On 27 April all three eggs hatched, giving an estimated incubation period of 24 days. Adults on the nest were often reluctant to flush, leaving only after observers approached within 1.5 m. Results of video observations made at the nest, detailing incubation rhythms and behaviours, will be reported elsewhere (RAG unpubl.).

The nest and eggs of Red-winged Wood-rail described here are similar to those reported for congeners, and the estimated incubation period of 24 days is also similar to that of other wood-rails⁴. The nest in the private Mushullacta community reserve appears to be the first confirmed report of

the species' presence within a formally protected area³. Breeding activity at nearly 1,200 m suggests that the elevational range given by Ridgely & Greenfield³ should be extended upwards by at least 300 m.

Acknowledgements

We are indebted to the community of Mushullacata and the Narvaez family, for their hospitality and access to their private reserve. This study was funded in part through the Population Biology Foundation via donations by John V. & the late Ruth Ann Moore, by a Rufford Small Grant, the Hertzberg Family Foundation, and a Pamela & Alexander F. Skutch Award. The PBNHS provided logistical and emotional support whilst in the field. Thanks to Murray Cooper for permitting use of his photograph. This is publication no. 61 of the Yanayacu Natural History Research Group.

References

1. BirdLife International (2004) *Threatened birds of the world 2004*. CD-ROM. Cambridge, UK: BirdLife International.
2. Greeney, H. F. (2005) *Aramides calopterus*, Red-winged Wood-rail. In: Greeney, H. F., Dobbs, R. C. & Martin, P. R. (eds.) *Natural history of Ecuador's mainland avifauna*. <http://depts.washington.edu/nhrp/nhema.htm>.
3. Ridgely, R. S. & Greenfield, P. J. (2001) *The birds of Ecuador*. Ithaca, NY: Cornell University Press.
4. Taylor, P. B. (1998) Family Rallidae (rails). In: del Hoyo, J., Elliott, A. & Sargatal, J. (eds.) *Handbook of the birds of the world*, 3. Barcelona: Lynx Edicions.

Jarol F. Vaca B. and Mike Tidwell

Shiripuno Research Center, Pastaza, Ecuador, c/o Joquin Pinto EI-353 y Amazonas, Quito, Ecuador. E-mail: vaca@shiripunoresearch.org.

Harold F. Greeney and Rudolphe A. Gelis

Yanayacu Biological Station & Center for Creative Studies, Cosanga, Napo, Ecuador, c/o 721 Foch y Amazonas, Quito, Ecuador.

Caroline Dingle

University of Cambridge, Zoology Department, Downing Street, Cambridge CB2 3EJ, UK.

Neils Krabbe

Zoological Museum, University of Copenhagen, Universitetsparken 15, DK-2100, Copenhagen, Denmark.